(12) INTERCATION CITY PER CATION PUBLISHED DESIGNATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



# . 1 (1881) 1 (1881) 1 (1881) 1 (1881) 1 (1881) 1 (1881) 1 (1881) 1 (1881) 1 (1881) 1 (1881) 1 (1881) 1 (1881) 1

(43) International Publication Date 17 July 2003 (17.07.2003)

**PCT** 

(10) International Publication Number WO 03/057203 A2

(51) International Patent Classification7:

\_\_\_

A61K 31/00

(21) International Application Number: PCT/EP03/00078

(22) International Filing Date: 7 January 2003 (07.01.2003)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

PD2002A000003 11 January 2002 (11.01.2002)

(71) Applicants (for all designated States except US): FIDIA FARMACEUTICI S.P.A. [IT/IT]; Via Ponte della Fabbrica, 3/A, I-35031 Abano Terme (IT). DEUTSCHES KREBSFORSCHUNGSZENTRUM (DKFZ) [DE/DE]; Im Neuenheimer Feld 280, 69120 Heidelberg (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): FUSENIG, Norbert, E. [DE/DE]; Am Pferchelhang 32, 69118 Heidelberg (DE). STARK, Hans-Jürgen [DE/DE]; Am Plötzberg 16, 74909 Meckesheim (DE). WILLHAUCK, Michael [DE/DE]; Rollbühlstrasse 58A, 68309 Mannheim (DE). PAVESIO, Alessandra [IT/IT]; Riviera Paleocapa, 9, I-35141 Padova (IT).

- (74) Agents: MINOJA, Fabrizio et al.; Bianchetti Bracco Minoja S.r.l., Via Rossini, 8, I-20122 Milano (IT).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

#### Published:

 without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

DEST AVAILABLE COPY

3/057203 A

(54) Title: BIOMATERIALS BASED ON HYALURONIC ACID FOR THE ANTI-ANGIOGENIC THERAPY IN THE TREATMENT OF TUMOURS

(57) Abstract: The use in the medical-surgical field of biomaterials based on hyaluronic acid derivatives, optionally in association with natural, synthetic or semisynthetic biopolymers, for suppressing the angiogenic process associated with tumour proliferation (in primary and secondary tumours) is disclosed.

WO 03/057203 PCT/EP03/00078

# BIOMATERIALS BASED ON HYALURONIC ACID FOR THE ANTI-ANGIOGENIC THERAPY IN THE TREATMENT OF TUMOURS

## SUBJECT OF THE INVENTION

5

10

15

20

25

The present invention relates to the use in the medical-surgical field of biomaterials based on hyaluronic acid derivatives, optionally in association with natural, synthetic or semisynthetic biopolymers, for suppressing the angiogenic process associated with tumour proliferation (in primary and secondary tumours).

## **BACKGROUND OF THE INVENTION**

The induction and development of angiogenesis is a pre-requisite for the development of a primary tumour, and for any subsequent metastases.

Angiogenesis is a dynamic process closely linked with the proliferation of cancer cells, because it is the latter that are chiefly responsible for the production and release of angiogenic factors, such as cytokines and other trophic factors. An increase in the vascularisation of a primary tumour can cause an increase in the number of cancer cells that enter into the circulation and give rise to new metastases.

Recent studies have demonstrated that an increase in the density of microvessels in an area affected by neoplasia indicates new tumour growth.

It is therefore clinically important to suppress angiogenesis to inhibit its development, if possible. Indeed, by associating anti-angiogenic therapy with "classic" anticancer therapy with drugs and/or radiation, with or without surgical removal of the tumour, it is possible to halt the proliferation of cancer cells, thus preventing the invasion of further tissues by said cells, and the consequent development of new metastases (Skobe H. et al., Nature Medicine, 1222-1227 (1997)).

In histological assessment of the onset of the angiogenic process

associated with a cancerous growth, it is important to look for markers of the tumour's vascular system, for example with antibodies that differentiate the endothelial cells from the cancerous ones. For example, the anti-CD3 antibody is specific for marking the endothelial cells and therefore enables their identification in the angiogenic process associated with the development of new metastases. Its use has proved essential in assessing the level of microvessel development associated with neoplasia. Indeed, thanks to antibody marking, it is possible to visualise and count the number of interconnections of the vessels within the cancerous tissue to understand and quantify the angiogenic process, relating it to any new developments in the neoplasia (thereby deciding if/how much/how to associate a therapy that modulates or inhibits angiogenesis with an established/classic anticancer therapy.

5

10

15

20

25

One such therapy consists in administering drugs that act by blocking the receptors of the trophic factors (PGDF, bFGF, VEGF) that are also angiogenic factors.

Preclinical results 'in vivo' have shown that said drugs prove important in inhibiting tumour growth but they do not determine regression of the tumour itself: on the strength of these major experimental data, the drugs have been introduced in numerous clinical trials.

However, an anti-angiogenic clinical therapy that provides for a generally oral pharmacological administration in chronic form may have many toxic side effects, because angiogenesis is not only associated with pathological disorders but also physiological processes such as tissue reproduction and repair ("Cancer: Principle Practice of Oncology" V. De Vita, S. Hellmann and S. Rosenberg, 6<sup>th</sup> Edition).

It is therefore of strategic importance to associate classic anticancer therapy with an anti-angiogenic therapy "in situ", and this is the subject of the

5

10

15

20

present invention.

Hyaluronic acid is one of the chief components of the extracellular matrix of the connective tissue, and there are numerous scientific publications concerning its role in various processes, both physiological and pathological, such as the formation of granulation tissue, chemotaxis in the inflammatory process, cell differentiation for various cell types. Other studies concern its role within the family of "substrate adhesion molecules".

Hyaluronic acid has been used for the above indications:

- as a differentiating agent in therapy for acute myeloid leukaemia (Charrad R. S. et al., Nature Medicine 5, 669-676 (1999));
- as a vehicle for drugs such as steroids or NSAIDs, antibiotics and anti-neoplastic agents, because of the abundant expression of its receptor (CD44) in cancer cells; (Freemantle, C. et al., Int. J. Tiss. Reac. XVIII (4) 157-166 (1995); Coradini, D. et al., Int. J. Cancer 5, 411-416 (1999));
- in preclinical studies on the inhibition of lung metastasis, because of its capacity for inhibiting the adhesion of cancer cells to the vascular endothelium (Karasaza K. et al., Clinical & Experimental Metastasis 15, 83-93 (1997));
- as a means of controlling adhesion to the substrate with subsequent proliferation of cells (possibly also cancer cells) permanently "in situ" after surgical removal of tissues (including tumours) (U.S. 5,627,162).

Experimental observations "in vivo" have, however, revealed that

25 hyaluronic acid may have a chemotaxic activity on cancer cells within the
granulation tissue that forms after removal of cutaneous metastasis of
melanoma (Salmon-Ehr, V. et al., Ann. Dermatol. Venereol, 123, 194-195
(1996)). Moreover, numerous pre-clinical studies have demonstrated that

hyaluronic acid enhances cancer cell migration, thereby favouring metastasis, as it is known that the degradation products of hyaluronic acid, that is, oligosaccharides constituted by 10 and 20 oligomers, are strong inducers of the angiogenic process (Hayen et al., J. Cell. Sci. 112, 2241-2251 (1999); Slevin, M. et al., Lab. Invest. 78(8), 987-1003 (1998)).

5

10

15

20

25

Moreover, biomaterials based on hyaluronic acid and/or the derivatives thereof have never been used as an anti-angiogenic therapy, neither have any other biodegradable and/or non-biodegradable biopolymers ever been used in anticancer therapies.

Absolutely innovative, therefore, is the use of biomaterials based on hyaluronic acid derivatives such as Hyaff® (EP 0 216 453 B1) or ACPs (EP 0 341 745 B1) in the form of non-woven felts for instance (EP0 618 817 B1) or as three-dimensional structures (WO 99/61080), possibly in association with various biomaterials (e.g. natural ones such as collagen, cellulose, polysaccharides, chitin, chitosan, pectin, agar, gellan and alginic acid, synthetic ones such as polylactic acid (PLA), polyglycolic acid (PGA), polyurethanes and polysulphonic resins, or semisynthetic ones such as collagen cross-linked with aldehyde, diamine and gellan) as a therapy to suppress and/or inhibit the angiogenic process that enhances and determines tumour metastasis.

## DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to biomaterials based on hyaluronic acid derivatives made into non-woven felts (as the preferred form of biomaterial), optionally in association with natural, synthetic or semisynthetic biopolymers, for use in the medical-surgical field as a new anti-angiogenic therapy ("in situ"), optionally associated with classic pharmacological anticancer therapies and/or radiotherapy, to modulate indirectly the proliferation of tumours, thus blocking the formation of local relapses and, therefore, any new metastases.

In order to study, characterise and then assess "in vivo" the effect of the biomaterial of the present invention in the angiogenic process that supports the development of skin carcinomas (considered to be a clarifying example), the Applicant has developed a new model of tumour/ stromal cell support interaction, described as follows:

5

10

15

20

- 1) two cell lines of human keratinocytes transfected with the rasoncogene: HACaT II-4, malignant variant and A5, benign variant;
- 2) said cells are transferred onto a collagen gel mounted into teflon rings covered by a silicone chamber, known as the Fusenig silicone chamber (FSC);
- 3) said FSC is then placed over the muscle fascia of the backs of nude mice, in the presence or absence of an immediately underlying layer of biomaterial based on Hyaff® 11 (total benzyl ester of hyaluronic acid) made in the form of a non-woven felt;
- 4) four to six weeks later, two different types of granulation tissue will have formed underneath the cancerous epithelium;
- 5) the development of the epithelial tumour and of the underlying granulation tissue is assessed, over time, both by classic histological analyses (haematoxylin/eosin) and by immunohistochemical techniques using the anti-CD31 antibody, to visualise the presence of vascular epithelium and therefore determine the development of the angiogenic process;
- 6) the levels of cellular proliferation are examined using immunohistochemical techniques associated with the introduction of BrdU into the DNA of proliferating cells, both within the granulation tissue underneath the epithelium and in the cancerous epithelium itself. Marking with the anti-integrin α6 antibody was also assessed to study the level of cellular proliferation within the

cancerous epithelium.

The results of the experiment were as follows:

## HACaT A5 line:

5

10

15

20

25

After 4-6 weeks, the cancerous epithelium in the control FSC (i.e. without any biomaterial placed under the epithelium), was well developed and multilayered, while the layer of granulation tissue underneath had completely replaced the layer of collagen that separated the epithelium from the underlying tissue (Fig. 1).

Conversely, four weeks later, the cancerous epithelium in the FSC placed over the Hyaff®-based biomaterial in the form of a non-woven felt is less developed than the relative control, and the layer of collagen that separates it from the nascent granulation tissue underneath is still thick and not infiltrated by cells and/or vessels (Fig. 1).

After six weeks, the quantity of collagen is still abundant, with just an initial layer of granulation tissue that begins to form over the biomaterial (Fig. 1).

## HACaT II 4 line:

After four to six weeks, in the control FSC, the cancer cells have constituted a thick epithelium that penetrates into the thickness of the new granulation tissue underneath, that has already completely replaced the layer of collagen that separated it from the epithelium (Fig. 2).

Four weeks later, in the FSC placed over the Hyaff®-based biomaterial, the cancerous epithelium is thin but easily distinguishable from the granulation tissue forming over the biomaterial, separated from this tissue by the collagen gel that is still present and not yet absorbed (Fig. 2).

Six weeks later, the tumour mass and the granulation tissue have established close contact, but there has been no actual infiltration of tumour cells into the granulation tissue, unlike the control, where the tumour cells have completely invaded the new, underlying granulation tissue (Fig. 2).

Using immunohistochemical techniques linked with the specific marking of particular nucleotides such as BrdU, at the 1<sup>st</sup> and 2<sup>nd</sup> weeks, good cell proliferation is evident within the nascent granulation tissue in the control and in the Hyaff®-based biomaterial, while at 4, and especially at 6, weeks after transplant, the cell growth rate drops drastically in the granulation tissue underneath the cancerous epithelium, which conversely maintains in both samples a good level of cellular proliferation (Fig. 3).

5

10

15

20

25

The growth of cancerous epithelium can also be visualised with a specific antibody against the protein integrin  $\alpha 6$ . Said molecule is, indeed, a component of the hemidesmosomes and its expression is normally only associated with the proliferative area of the epithelial layers.

Fig. 3a shows that antibody marking for the integrin protein α6 is notably present throughout the cancerous epithelium both in the control FCS and in the FCS with the Hyaff®-based biomaterial, even though expression of the test protein appears less extensive throughout the thickness of the cancerous epithelium in the latter sample.

Specific marking for the vascular epithelium with the anti-CD31 antibody reveals, furthermore, that in the controls, after four weeks, the angiogenic process is well established as the vessels in the underlying granulation tissue already reach the cancerous epithelium and after 6 weeks they invade it, thus favouring metastasis (Fig. 4).

In the case of the FSC with the Hyaff®-based biomaterial, after four weeks there is still no close contact between granulation tissue and cancerous epithelium. This will occur only after six weeks, even though there is no invasion of the epithelium by the underlying microvessels, that remain relegated to the granulation tissue (Fig. 4).

The angiogenic process seems to be at a standstill, no longer enhancing

tumour development. Vascularisation is limited to the area covered by the Hyaff®-based biomaterial, so the tumour cells do not invade the granulation tissue that has formed within the biomaterial.

In conclusion, the Hyaff®-based biomaterial proved able to modulate/inhibit the angiogenic process related to vascularisation of the cancerous epithelium. It therefore proves to be particularly advantageous to use the biomaterials based on hyaluronic acid derivatives in the oncological field, where it is important to modulate the angiogenic process and therefore, indirectly, the proliferation of cancer cells in primary and secondary tumours.

5

10

15

20

25

According to the invention, the biomaterials that can be useful in the oncological field as a new anti-angiogenic therapy "in situ" may be, for example, in the form of non-woven felts, sponges, films, membranes, microspheres or in other three-dimensional forms in cases where it is necessary to fill the cavities that are liable to form after surgical removal of a tumour.

The anti-angiogenic action of the biomaterial can, moreover, be supported by supplementing the biomaterial with NSAIDs, steroids, hormones, antibiotics and especially anti-cancer drugs such as fluorouracil, methotrexate, cis-platinum, carboplatin, oxaliplatin, ethopoxide, cyclophosphamide, vincristine, doxorubicin.

The invention being thus described, it is clear that these methods can be modified in various ways. Said modifications are not to be considered as divergences from the spirit and purposes of the invention and any modification that would appear evident to an expert in the field comes within the scope of the following claims.

## **CLAIMS**

5

- 1. Biomaterials constituted by at least one hyaluronic acid derivative, optionally in association with other natural, synthetic and/or semisynthetic biopolymers and with pharmacologically active substances, as an anti-angiogenic therapy to treat primary and secondary tumours.
- 2. Biomaterials according to claim 1, wherein the hyaluronic acid derivative is a benzyl ester.
- 3. Biomaterials according to claim 1, wherein the hyaluronic acid derivative is cross-linked.
  - 4. Biomaterials according to claim 1, wherein the natural biopolymer is selected from the group consisting of collagen, cellulose, polysaccharides, chitin, chitosan, pectins, agar, gellan and alginic acid.
  - 5. Biomaterials according to claim 1, wherein the synthetic biopolymer is selected from the group consisting of polylactic acid (PLA), polyglycolic acid (PGA), polyurethanes and polysulphonic resins.
  - 6. Biomaterials according to claim 1, wherein the semisynthetic biopolymer is selected from the group consisting of collagen cross-linked with aldehydes, diamine and gellan.
- 7. Biomaterials according to claim 1, wherein the biopolymer may optionally be in association with pharmacologically active substances such as fluorouracil, methotrexate, cis-platinum, carboplatin, oxaliplatin, ethopoxide, cyclophosphamide, vincristine, doxorubicin.
- 8. The use of biomaterials constituted by at least one hyaluronic acid derivative, optionally in association with other natural, synthetic and/or semisynthetic biopolymers and with pharmacologically active substances, as an anti-angiogenic therapy for the treatment of primary and secondary tumours.

- 9. The use of biomaterials according to claim 8, wherein the hyaluronic acid derivative is a benzyl ester.
- 10. The use of biomaterials according to claim 8 wherein the hyaluronic acid derivative is cross-linked.
- 5 11. The use of biomaterials according to claim 8 wherein the natural biopolymer is selected from the group consisting of collagen, cellulose, polysaccharides, chitin, chitosan, pectins, agar, gellan and alginic acid.
  - 12. The use of biomaterials according to claim 8 wherein the synthetic biopolymer is selected from the group consisting of polylactic acid (PLA), polyglycolic acid (PGA), polyurethanes and polysulphonic resins.

10

- 13. The use of biomaterials according to claim 8 wherein the semisynthetic biopolymer is collagen cross-linked with aldehydes, diamine and gellan.
- 14. The use of biomaterials according to claim 8 wherein the biopolymer may optionally be in association with pharmacologically active substances such as fluorouracil, methotrexate, cis-platinum, carboplatin, oxaliplatin, ethopoxide, cyclophosphamide, vincristine and doxorubicin.
- 15. The use of a biomaterial according to claim 8 wherein the hyaluronic acid derivative is made into a non-woven felt, sponge, microsphere, film, membrane and/or other three-dimensional structures.
- 20 16. The use of biomaterials constituted by at least one hyaluronic acid derivative, optionally in association with other natural, synthetic and/or semisynthetic biopolymers and with pharmacologically active substances, for the treatment and care of primary and secondary tumours when the tumour has been surgically removed and the cavity that is thus formed requires filling, making it advantageous to prevent and/or inhibit the angiogenic process using the biomaterial itself.

1/5




Fig. 1

2/5




Fig. 2





Fig. 3
SUBSTITUTE SHEET (RULE 26)



5/5




Fig. 4

#### (19) World Intellectual Property **Organization** International Bureau





(43) International Publication Date 17 July 2003 (17.07.2003)

(10) International Publication Number WO 2003/057203 A3

(51) International Patent Classification<sup>7</sup>: A61P 35/00, A61K 47/36

A61L 27/00,

(74) Agents: MINOJA, Fabrizio et al.; Bianchetti Bracco Minoja S.r.l., Via Rossini, 8, I-20122 Milano (IT).

(21) International Application Number:

PCT/EP2003/000078

(22) International Filing Date: 7 January 2003 (07.01.2003)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

PD2002A000003

11 January 2002 (11.01.2002) IT

- (71) Applicants (for all designated States except US): FIDIA FARMACEUTICI S.P.A. [IT/IT]; Via Ponte della Fabbrica, 3/A, I-35031 Abano Terme (IT). DEUTSCHES KREBSFORSCHUNGSZENTRUM (DKFZ) [DE/DE]; Im Neuenheimer Feld 280, 69120 Heidelberg (DE).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): FUSENIG, Norbert, E. [DE/DE]; Am Pferchelhang 32, 69118 Heidelberg (DE). STARK, Hans-Jürgen [DE/DE]; Am Plötzberg 16, 74909 Meckesheim (DE). WILLHAUCK, Michael [DE/DE]; Rollbühlstrasse 58A, 68309 Mannheim (DE). PAVESIO, Alessandra [IT/IT]; Riviera Paleocapa, 9, I-35141 Padova (IT).

- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
- LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

#### Published:


- with international search report
- (88) Date of publication of the international search report: 31 December 2003

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: BIOMATERIALS BASED ON HYALURONIC ACID FOR THE ANTI-ANGIOGENIC THERAPY IN THE TREAT-MENT OF TUMOURS

(57) Abstract: The use in the medical-surgical field of biomaterials based on hyaluronic acid derivatives, optionally in association with natural, synthetic or semisynthetic biopolymers, for suppressing the angiogenic process associated with tumour proliferation (in primary and secondary tumours) is disclosed.





A. CLASSIFICATION OF SUBJECT MATTER IPC 7 A61L27/00 A61P35/00 A61K47/36

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols) IPC 7 A61K A61L

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, MEDLINE, BIOSIS, WPI Data, PAJ

	ENTS CONSIDERED TO BE RELEVANT	Relevant to claim No.
Category *	Citation of document, with indication, where appropriate, of the relevant passages	
P,X	WO 02 18448 A (FRANCESCANGELI ANDREA;BELLINI DAVIDE (IT); CRESCENZI VITTORIO (IT) 7 March 2002 (2002-03-07) page 3, line 9-12 claims 5,11,15,24	1–16
P,X	WO 02 41877 A (CLEAR SOLUTIONS BIOTECH INC; DEHAZYA PHILIP (US); LU CHENG (US)) 30 May 2002 (2002-05-30) page 4, line 19-22 page 18, line 8 page 16, line 21	1-16
P,X	WO 02 18450 A (FRANCESCANGELI ANDREA; RENIER DAVIDE (IT); CRESCENZI VITTORIO (IT)) 7 March 2002 (2002-03-07) page 2, line 10-12 page 10, line 27	1-16

Further documents are listed in the continuation of box C.	Patent family members are listed in annex.
Special categories of cited documents:  A document defining the general state of the art which is not considered to be of particular relevance  E earlier document but published on or after the international filing date  L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)  O document referring to an oral disclosure, use, exhibition or other means  P document published prior to the international filing date but later than the priority date claimed	<ul> <li>'T' later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</li> <li>'X' document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to the involve an inventive step when the document is taken alone</li> <li>'Y' document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</li> <li>'&amp;' document member of the same patent family</li> </ul>
Date of the actual completion of the international search	Date of mailing of the international search report
26 June 2003	10/07/2003
Name and mailing address of the ISA European Palent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk	Authorized officer
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Heller, D

Int Interior Application No
PCT/EP 03/00078

	otion) DOCUMENTS CONSIDERED TO BE RELEVANT  Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Category °	Citation of document, with indication, where appropriate, of the following passages	
х	EP 0 341 745 A (FIDIA SPA) 15 November 1989 (1989-11-15) page 1, line 32 page 5, line 41-49 page 6, line 19-28 page 9, line 55 page 10, line 11 page 10, line 14	1-7
X	WO 99 61080 A (CALLEGARO LANFRANCO ;DONA MASSIMO (IT); FIDIA ADVANCED BIOPOLYMERS) 2 December 1999 (1999-12-02) page 1, line 6-13 page 4, line 23-28	1-7
X	US 5 520 916 A (CALLEGARO LANFRANCO ET AL) 28 May 1996 (1996-05-28) column 1, line 8-13 claim ALL	1-7
X	WO 00 01733 A (BELLINI DAVIDE ;FIDIA ADVANCED BIOPOLYMERS SRL (IT); TOPAI ALESSAN) 13 January 2000 (2000-01-13) page 1, line 11-14 claims 1,5,16,21,23	1-16
X	WO 00 54762 A (UNIV BOSTON ; MOULTON STEVEN (US)) 21 September 2000 (2000-09-21) claims 1-31	1-16
X	WO 00 57896 A (MUMMERT MARK E ;TAKASHIMA AKIRA (US); UNIV TEXAS (US); MOHAMADZADE) 5 October 2000 (2000-10-05) page 3, line 21-32	1-16
X	US 4 851 521 A (DELLA VALLE FRANCESCO ET AL) 25 July 1989 (1989-07-25) column 1, line 6-12 claims 1-8,25	1-16
X	EP 0 466 300 A (BIOMATRIX INC) 15 January 1992 (1992-01-15) page 3, line 35-47 page 7, line 28-31	1-16
P,X	TONELLO C ET AL: "In vitro reconstruction of human dermal equivalent enriched with endothelial cells" BIOMATERIALS, ELSEVIER SCIENCE PUBLISHERS BV., BARKING, GB, vol. 24, no. 7, March 2003 (2003-03), pages 1205-1211, XP004401497 ISSN: 0142-9612 abstract	1-7
1	-/	<b>\</b>

Intermional Application No
PCT/EP 03/00078

C.(Continu	ation) DOCUMENTS CONSIDERED TO BE RELEVANT	Relevant to claim No.
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Helevani to Claim No.
X	GLASS J ET AL: "A three-dimensional cell attachment matrix created by cross-linking RGD peptide modified hyaluronic acid" JOURNAL OF CELLULAR BIOCHEMISTRY - SUPPLEMENT, WILEY-LISS, US, no. SUPPL 19A, 5 January 1995 (1995-01-05), page 178 XP002112805 ISSN: 0730-2312 abstract	1-7
X	LUO Y ET AL: "Cross-linked hyaluronic acid hydrogel films: new biomaterials for drug delivery" JOURNAL OF CONTROLLED RELEASE, ELSEVIER SCIENCE PUBLISHERS B.V. AMSTERDAM, NL, vol. 69, no. 1, 3 October 2000 (2000-10-03), pages 169-184, XP004217543 ISSN: 0168-3659 abstract	1-7
X	CORADINI D ET AL: "Hyaluronic acid as drug delivery for sodium butyrate: improvement of the anti-proliferative activity on a breast-cancer cell line." INTERNATIONAL JOURNAL OF CANCER. JOURNAL INTERNATIONAL DU CANCER. UNITED STATES 5 MAY 1999, vol. 81, no. 3, 5 May 1999 (1999-05-05), pages 411-416, XP002245583 ISSN: 0020-7136 abstract	1-16

Intermonal Application No
PCT/EP 03/00078

				1017	L1 05/00070
Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 0218448	A	07-03-2002	ΙT	PD20000208 A1	28-02-2002
WU 0210440	^	07 03 2002	ĀÙ	9181501 A	13-03-2002
			WO	0218448 A2	07-03-2002
					02.06.2002
WO 0241877	Α	30-05-2002	AU	3969702 A	03-06-2002
			WO	0241877 A1	30-05-2002
			US	2003096734 A1	22-05-2003
UO 0219450	Α	07-03-2002	IT	PD20000207 A1	28-02-2002
WO 0218450	A	07 03 2002	ĀÚ	1387002 A	13-03-2002
			WO	0218450 A1	07-03-2002
			ËΡ	1313772 A1	28-05-2003
					10.05.1000
EP 0341745	Α	15-11-1989	IŢ	1219587 B	18-05-1990 15-12-1994
			AT	115590 T	15-12-1994
			AT	195534 T	
			ΑU	631125 B2	19-11-1992 29-11-1989
			AU	3574789 A	29-11-1989 29-07-1997
			CA	1339122 C	29-07-1997 26-01 <b>-</b> 1995
			DE	68919900 D1	11-05-1995
			DE	68919900 T2	21-09-2000
			DE	68929241 D1	05-04-2001
			DE	68929241 T2	12-03-1990
			DK	10990 A	16-11-1989
			MO	8910941 A1	15-11-1989
			EP	0341745 A1	14-09-1994
			EP	0614914 A2 2064378 T3	01-02-1995
			ES		16-01-2001
			ES	2151910 T3 107050 B1	31-05-2001
			FI	3015035 T3	31-05-1995
			GR	3015035 T3 3034651 T3	31-01-2001
			GR	53666 A2	28-11-1990
			HU	90274 A	12-09-1996
			IL JP	10324701 A	08-12-1998
			JP	2504163 T	29-11-1990
			JP	2941324 B2	25-08-1999
			NZ	229100 A	28-08-1995
			US	5676964 A	14-10-1997
WO 9961080	Α	02-12-1999	IT	PD980131 A1	
			ΙT	1302535 B1	05-09-2000
			AT	227589 T	15-11-2002
			AU	748303 B2	30-05-2002
			AU	4368099 A	13-12-1999
			CA	2332532 A1	
			DE	69903945 D1	
			DK	1085917 T3	
			WO	9961080 A1	
			EP	1085917 A1	
			ES	2184460 T3	01-04-2003 04-06-2002
		•		2002616167	
			JP	2002516154 T	
 us 5520916	A			1254704 B	09-10-1995
 US 5520916	A	28-05-1996	JP	1254704 B 200630 T	09-10-1995 15-05-2001
 US 5520916	Α	28-05-1996	JP IT	1254704 B 200630 T 669147 B2	09-10-1995 15-05-2001 30-05-1996
 US 5520916	Α	28-05-1996	JP IT AT	1254704 B 200630 T	09-10-1995 15-05-2001

Interional Application No
PCT/EP 03/00078

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
US 5520916			BG	302 Y1	31-05-1999
07 DOZUATO	^		CA	2126085 A1	24-06-1993
			DE	69231796 D1	23-05-2001
			DE	69231796 T2	15-11-2001
			DK	618817 T3	30-07-2001
			WO	9311803 A1	24-06-1993
			EP	0618817 A1	12-10-1994
			ES	2155832 T3	01-06-2001
			FI	942894 A	18-08-1994
			GR	3036197 T3	31-10-2001
			ΗÜ	68680 A2	28-07-1995
			JP	7502430 T	16-03-1995
			NO	942330 A	17-08-1994
			NZ	246575 A	24-04-1997
			PT	618817 T	30-08-2001
			RO	115017 B1	29-10-1999
			RU	2133127 C1	20-07-1999
			US	5824335 A	20-10-1998
WO 0001733	Α	13-01-2000	ΙT	PD980169 A1	07-01-2000
			AU	4639799 A	24-01-2000
			CA	2339066 A1	13-01-2000
			EP	1095064 A1	02-05-2001
			WO	0001733 A1	13-01-2000
			JP	2002519481 T	02-07-2002
	Λ	21-09-2000	AU	3886600 A	04-10-2000
WO 0054762	Α	21-09-2000	CA	2365767 A1	21-09-2000
			EP	1162984 A2	19-12-2001
			JP	2002539157 T	19-11-2002
			WO	0054762 A2	21-09-2000
			US	2002169144 A1	14-11-2002
			US	6472379 B1	29-10-2002
WO 0057896	Α	05-10-2000	AU	4025500 A	16-10-2000
			CA	2366604 A1	05-10-2000
			EP	1165112 A1	02-01-2002
			JP	2002540161 T	26-11-2002
			WO	0057896 A1	05-10-2000
			US	2003054991 A1	20-03-2003
US 4851521	A	25-07-1989	IT	1203815 B	23-02-1989
03 4031321	• • • • • • • • • • • • • • • • • • • •	20 0, 2000	ĪŤ	1214658 B	18-01-1990
			ĀŤ	135713 T	15-04-1996
			AT	227741 T	15-11-2002
			ΑÙ	591501 B2	07-12-1989
			AU	5983686 A	26-02-1987
			CA	1341276 C	31-07-2001
			DE	3650501 D1	25-04-1996
			DE	3650501 T2	21-11-1996
			DE	3650776 D1	19-12-2002
			DK	323686 A	09-01-1987
			EP	0216453 A2	01-04-1987
			EP	0696598 A1	14-02-1996
					01-06-1988
			£ζ	2001512 An	01_00_1200
			ES	2001512 A6 862878 A .B.	
		·	ES FI FI	2001512 A6 862878 A ,B, 892710 A ,B,	01-00-1988 09-01-1987 02-06-1989

Intermonal Application No
PCT/EP 03/00078

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
US 4851521	A		FI	94767 B	14-07-1995
JU , UU			HU	42512 A2	28-07-1987
			ΙE	81120 B1	22-03-2000
			IL	79362 A	31-07-1995
			IN	165582 A1	25-11-1989
			JP	2569012 B2	08-01-1997
			JP	62064802 A	23-03-1987
			KR	8701901 B1	21-10-1987
			NO	862734 A ,B,	09-01-1987
			NO	910295 A	09-01-1987
			NZ	216786 A	26-05-1992
			NZ	233045 A	26-05-1992
			PH	25189 A	27-03-1991
			PT	82941 A ,B	01-08-1986
			SG	90006 A1	23-07-2002
			ÜS	4965353 A	23-10-1990
			US	5202431 A	13-04-1993
			US	5336767 A	09-08-1994
			PH	25383 A	03-06-1991
			PH	25382 A	03-06-1991
			PH	25642 A	21-08-1991
			PH	25643 A	21-08-1991
			ZA	8605071 A	30-09-1987
EP 0466300	Α	15-01-1992	US	5143724 A	01-09-1992
E1 0400300	,,	20 02 2002	AT	165979 T	15-05-1998
			AU	629467 B2	01-10-1992
			AU	7405591 A	09-01-1992
			CA	2041074 A1	10-01-1992
			DE	69129391 D1	18-06-1998
			DE	69129391 T2	08-10-1998
			DK	466300 T3	07-10-1998
			EP	0466300 A2	15-01-1992
			ĒS.	2117634 T3	16-08-1998
			HK	1011502 A1	09-07-1999
			JP	4261664 A	17-09-1992
			JP	7093943 B	11-10-1995
			US	5399351 A	21-03-1995
			ÜŠ	5246698 A	21-09-1993

# This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

□ BLACK BORDERS
□ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
□ FADED TEXT OR DRAWING
□ BLURRED OR ILLEGIBLE TEXT OR DRAWING
□ SKEWED/SLANTED IMAGES
□ COLOR OR BLACK AND WHITE PHOTOGRAPHS
□ GRAY SCALE DOCUMENTS
□ LINES OR MARKS ON ORIGINAL DOCUMENT
□ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

## IMAGES ARE BEST AVAILABLE COPY.

OTHER:

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.